


Abstract

We aim to use synthetic biology to engineer pro-biotic lactic acid bacteria used in the production of dairy products like yogurt, buttermilk and curds, to produce Monellin, a heat and pH stable sweetening protein. We plan to use the decreasing pH during curdling and the addition of nisin as the two inputs, and the CRE gene in combination with loxP sites as an AND gate. In addition, we have used the unique functionality of CRE to create a system that is self-excising and will return the GMO to 'wild-type' once the production is complete. We have managed to identify and characterise a pH-sensitive promoter, P170 and have abstracted SP310mut2 and CRE to be used in our system.

Concept

To conceptualise and synthesise a regulatory mechanism that functions as a pseudo-AND gate in response to two inputs.

Our pseudo-AND system is based on the gene system known as CRE-lox. We came across the CRE-Lox system, originally discovered in mammalian cells, but very effectively transferred to bacterial systems as a recombinase. Specifically, Cre-Lox recombination is a special type of site-specific recombination, which involves the targeting of a specific sequence of DNA and splicing it with the help of an enzyme called Cre recombinase.



It can be seen that only one of the combinations of inputs give rise to an output, simulating a pseudo-AND gate. It is important to note that the Gene-of-interest is excised permanently, returning the cell (and its daughters) to 'wild-type'.

References



- Morris J A, et al; Purification of monellin, the sweet principle of *D.cumminsii*, *Biochem Biophys Acta* (1972), vol 261:114
 Madsen S M, et al; Molecular characterisation of the pH-inducible and growth phase-dependent promoter P170 of *L.lactis*; *Molecular Microbiology* (1999), vol 32, pg 75 - 87
 Ravn P, et al; Optimisation of signal peptide SP310 for heterologous protein production in *L.lactis*; *Microbiology* (2003), vol 149, pg 2193 - 2201
 Pfeifer A, et al; Delivery of the Cre recombinase by a self-deleting lentiviral vector: efficient gene targeting in vivo.

For further information and contact details of the IIT_Madras team, please visit our wiki at http://2010.igem.org/Team:IIT_Madras.

Probiotic Sweetener, controlled by a pseudo-AND gate

Team: IIT_Madras


iGEM 2010 Jamboree



Construct Design


NICE

The Nisin-inducible expression system is upregulated under the presence of food-grade nisin in the system. It is also to be noted that the system should only be activated after the culture reaches an OD600 of around 0.7. The system is tested with a reporter with J23119 as a reference.



SP310mut2

The export tag is that which is obtained from the same vector as P170 and is an optimised version of the naturally found SP310. The system is tested under J23119 with a reporter with the same reporter without the export tag as a reference.




CRE-loxP

The CRE enzyme is a recombinase that modifies / excises the DNA fragment that lies between its specific recognition sites, named lox. We have chosen to test the system with two reporters, and the promoter J23119.



P170

This is a pH-sensitive growth phase-dependent promoter that upregulates expression from a baseline between a pH of 6.0 and 6.5. The system is to be tested with a reporter with J23119 as a reference.




Legend

- loxP
- Terminator
- RBS
- Coding Sequence
- Promoter


Results

P170

The testing of P170 was done at 5 different pH (4.5, 5, 5.5, 6, 7) with Firefly luciferase from pGL3Basic as the reporter. The data, given in Relative Luminescence Units/sec was normalised with OD600 and plotted against pH. The following trend was observed in the case of expression under P170.





To our surprise, we also discovered that the constitutive promoter, which we used as a reference promoter showed up-regulation at low pH (4.5 - 5) indicating that it might be under the influence of a pH-based stress factor.




Validation

CRE

To test the expected behavior of the CRE system, a plate based experiment is setup with the following construct and two reporters. On the transformation plate, different colonies will express the reporters based on the activity of the CRE within them. If the CRE functions within a particular cell, none of its future generations will carry the excised DNA fragment, therefore reporter 2 is expressed. Now, if CRE doesn't function upto a certain generation, reporter 1 is expressed and the terminator stops the expression of reporter 2. If the reporters can be visualised independently then it may be possible to identify the point at which excision occurred.



Option 1
CRE is not expressed,
Reporter 1 is expressed.



Option 2
CRE is expressed,
Reporter 1 is excised,
Reporter 2 is expressed.

SP310mut2

The characterisation for the export tag will also be done in pGL3Basic with Firefly luciferase as a reporter. Samples will be taken over time and the luciferase will be assayed for both before and after cell lysis, to estimate export efficiency. Luminescence measurements of the luciferase were made in conjunction with added luciferin. The colonies are ready and the experiment will be conducted once we return from the Jamboree.

Our Team

Undergraduates:

- | | | |
|-----------------------|---------------------|---------------------|
| Nishant | Avaneesh T | Sonia Chothani |
| Harsh Nagpal | Chaitra Prabhakar | Varun Chokshi |
| Srivathsan Adivarahan | Kousik Sundararajan | Akshay Rangamani |
| Sriram Srikant | Mahak Singhal | Chetan Srinath |
| Amarnath Bhat M | Praneet M | Vivek Parthasarathy |

Advisors and Instructors:

- Dr Guhan Jayaraman
 Dr K Chandraraj
 Shashi Bala Prasad
 Parashuraman Sohnewale

Our Sponsors



We also acknowledge the help of:

